

First and Last Name: \_\_\_\_\_

2. **Correct The Mistake**

Determine whether these equations are false, and if so write the correct answer.

(a)  $\ln\left(\frac{x}{2}\right) + \ln\left(\frac{x}{2}\right) = \ln(x)$

(b)  $e^x e^y = e^{xy}$

(c)  $\frac{d}{dx} \cos(4x^2) = \sin(4x^2)$

(d)  $\int_0^\infty dx x e^{-x^2} = \infty$

**Solution:**

- (a) This is an incorrect application of the rule for adding logarithms. When we add two logarithms with the same base, we obtain a new logarithm whose argument is a product of the arguments of the original two logarithms. The correct calculation is

$$\ln\left(\frac{x}{2}\right) + \ln\left(\frac{x}{2}\right) = \ln\left(\frac{x^2}{2}\right). \quad (1)$$

- (b) This is an incorrect application of the rule for multiplying exponentials. When we multiply two logarithms, we obtain a new exponential whose argument is a sum of the arguments of the original two exponentials. The correct calculation is

$$e^x e^y = e^{x+y}. \quad (2)$$

- (c) This calculation has the incorrect derivative for  $\cos(u)$  and does not apply the chain rule. The correct calculation is

$$\frac{d}{dx} \cos(4x^2) = -8x \sin(4x^2). \quad (3)$$

- (d) A definite integral over an infinite domain is not necessarily infinite. Instead, we need to use  $u$ -substitution to evaluate this integral. Given that  $\frac{d}{dx} e^{-x^2}/2 = -x e^{-x^2}$ , we have

$$\int_0^\infty dx x e^{-x^2} = -\frac{e^{-x^2}}{2} \Big|_0^\infty = \frac{1}{2}. \quad (4)$$

■

First and Last Name: \_\_\_\_\_

### 3. Probability

We have a large number of atoms in a system. Each atom decays at a rate  $\lambda$  such that after a time  $t$ , there is the probability  $\lambda t$  for the atom to have decayed. The probability that  $n$  atoms have decayed in time  $t$  is

$$p_n = \frac{(\lambda t)^n}{n!} e^{-\lambda t}, \quad (5)$$

where  $n! = n(n-1) \cdots 2 \cdot 1$ . What is  $\langle n \rangle$ , the average number of atoms that have decayed in a time  $t$ ?  
*Hint: The Taylor series of  $e^x$  is  $e^x = \sum_{n=0}^{\infty} x^n/n!$ .*

#### Solution:

By the definition of average, we find that the average number of atoms that have decayed in a time  $t$  is

$$\begin{aligned} \langle n \rangle &= \sum_{n=0}^{\infty} n p_n \\ &= \sum_{n=0}^{\infty} n \frac{(\lambda t)^n}{n!} e^{-\lambda t} \\ &= \lambda t \sum_{n=1}^{\infty} \frac{(\lambda t)^{n-1}}{(n-1)!} e^{-\lambda t} \\ &= \lambda t e^{-\lambda t} \sum_{k=0}^{\infty} \frac{(\lambda t)^k}{k!}. \end{aligned} \quad (6)$$

In the third equality, we factored  $\lambda t$  from the expression, used  $n/(n!) = 1/(n-1)!$ , and began our summation from  $n = 1$  because the  $n = 0$  term vanishes. Using the Taylor series for the exponential, we then find

$$\langle n \rangle = \lambda t e^{-\lambda t} e^{\lambda t} = \lambda t. \quad (7)$$

■

First and Last Name: \_\_\_\_\_

4. **Probability, II**

We have a random number generator in which each number from 0 to 999 is equally likely to occur. What is the probability of getting 547?

We change the random number generator so that there is a probability of  $1/2$  of getting a 9 in the hundreds digit and a probability of  $1/2 \times 1/9$  of getting any particular other number (e.g., a probability of  $1/18$  of getting a 2) in the hundreds digit. The probability distribution for the tens and ones digits remain unchanged. What is the probability of getting 547?

**Solution:**

If we have a random number generator in which each number from 0 to 999 is equally likely to occur, then the probability of getting 547 is the inverse of the total number of possible numbers. Namely,

$$p_{547} = \frac{1}{1000}. \quad (8)$$

If we have a random number generator in which the probability of getting a 9 in the hundreds digit is  $1/2$  and the probability of getting any other particular number is  $1/2 \times 1/9 = 1/18$ , (with the probability distribution of getting other numbers unchanged), the probability of obtaining 547 is

$$p_{547} = \frac{1}{18} \times \frac{1}{10} \times \frac{1}{10} = \frac{1}{1800}. \quad (9)$$

■

First and Last Name: \_\_\_\_\_

### 5. Information

Our friend performs some random trial which yields a random number. We define "information" needed to specify the number as the average number of binary-valued questions that need to be answered to determine the hidden number. Determine the "information" needed to specify the number given that our friend conducted the following random trials

- (a) Selecting a card from a deck of cards with numbers from 1 to 100 (each number occurring once)
- (b) Selecting a card from a deck of cards with numbers from 1 to 100 where even numbers are twice as likely as odd numbers.

### Solution:

- (a) In Lecture Notes 03 "Entropy and Information", we determined that if there is a probability  $p_i$  of getting a number  $i$  out of a set of numbers  $\{i\}$ , then the average number of questions we will need to guess the number (under the given conditions of the "Guess that number" game) is

$$\langle \# \text{ of Qs} \rangle = - \sum_{\{i\}} p_i \log_2 p_i. \quad (10)$$

If we have a deck of cards numbered 1 to 100, and we select one card, there is a uniform probability of  $1/100$  for getting any particular card. Thus the amount of information needed to specify the selected card is

$$\langle \# \text{ of Qs} \rangle = \log_2(100) \approx 6.64. \quad (11)$$

- (b) If even numbers are twice as likely as odd numbers, then the probability of getting an even number is  $2/3$  and the probability of getting an odd number is  $1/3$ . Since there are 50 even numbers and 50 odd numbers for a deck of cards from 1 to 100, the probability of getting any particular even number is  $2/3 \times 1/50 = 1/75$  and the probability of getting any particular odd number is  $1/3 \times 1/50 = 1/150$ . From these probabilities, we find that the amount of information needed to specify a drawn card from this deck is

$$\begin{aligned} \langle \# \text{ of Qs} \rangle &= - \sum_{\{i\}} p_i \log_2 p_i \\ &= - \sum_{\text{even \#s}} \frac{1}{75} \log_2 \frac{1}{75} - \sum_{\text{odd \#s}} \frac{1}{150} \log_2 \frac{1}{150} \\ &= - \frac{50}{75} \log_2 \frac{1}{75} - \frac{50}{150} \log_2 \frac{1}{150} \\ &= \frac{2}{3} \log_2 75 + \frac{1}{3} \log_2 150 \\ &= \frac{1}{3} \log_2 (75^2 \times 150) \approx 6.56. \end{aligned} \quad (12)$$

■

First and Last Name: \_\_\_\_\_

**6. Entropy of a Macrostate**

We have a spin system with  $N$  spins of which  $n_{\uparrow}$  spins are pointing up and  $N - n_{\uparrow}$  spins are pointing down. What is the entropy of the macrostate characterized by a given  $n_{\uparrow}$ , given that each associated microstate is equally likely? *Hint: Entropy is  $k_B$  times the natural logarithm of the number of microstates for a particular macrostate.*

**Solution:**

For a spin system with  $N$  spins of which  $n_{\uparrow}$  spins are pointing up and  $N - n_{\uparrow}$  spins are pointing down, then the number of microstates associated with this macrostate is

$$\Omega = \binom{N}{n_{\uparrow}}. \quad (13)$$

Each of the microstates with a common  $N$  and  $n_{\uparrow}$  are equally likely, so the entropy of the macrostate (or set of microstates) with  $N$  and  $n_{\uparrow}$  is

$$S = k_B \ln \Omega = k_B \ln \binom{N}{n_{\uparrow}}. \quad (14)$$

■

First and Last Name: \_\_\_\_\_

**6. Maximization with constraint**

The perimeter of a rectangle with side lengths  $x$  and  $y$  is given  $P = 2x + 2y$ . The area of the rectangle is given by  $A = xy$ . In terms, of  $P$  for what value of  $x$  is  $A$  maximized? Given this  $x$ , what is  $y$ ? *Note: You might be able to guess the answer but I also want you to show it analytically.*

**Solution:**

We will find the maximum area by writing  $A$  exclusively as a function of one of the sides of the rectangle and then implementing the local maximization algorithm. First, we note that if  $P = 2x + 2y$ , then

$$y = \frac{1}{2}(P - 2x), \tag{15}$$

where  $P$  is the perimeter of the rectangle. The area as an exclusive function of  $A$  is then

$$A = xy = \frac{x}{2}(P - 2x). \tag{16}$$

We are now, looking for the value of  $x$ , which maximizes  $A$ . Namely, the  $x$  such that  $A'(x) = 0$  but  $A''(x) < 0$ . First computing  $A'(x)$ , we have

$$A'(x) = \frac{P}{2} - 2x. \tag{17}$$

Setting  $A'(x) = 0$ , we find that  $A(x)$  *might* have a maximum at  $x = P/4$  and (by Eq.(15))  $y = P/4$ . We affirm the local maximum condition by noting that

$$A''(x) = -2 < 0, \tag{18}$$

Therefore,  $x = y = P/4$ , subject to the constraint  $P = 2x + 2y$ , indeed maximizes the area  $A = xy$ . ■

First and Last Name: \_\_\_\_\_

7. **Gaussian integral**

Compute the integral

$$\int_{-\infty}^{\infty} dx_1 \int_{-\infty}^{\infty} dx_2 \cdots \int_{-\infty}^{\infty} dx_N \exp\left(-\sum_{i=1}^N \lambda_i x_i^2\right).$$

Write the final result using the product symbol  $\prod$ .

**Solution:**

To evaluate the given multi-dimensional integral, we first recall the identity

$$\int_{-\infty}^{\infty} dx e^{-\lambda x^2} = \sqrt{\frac{\pi}{\lambda}}. \quad (19)$$

Now, turning the the multi-dimensional integral, we have

$$\begin{aligned} \int_{-\infty}^{\infty} dx_1 \int_{-\infty}^{\infty} dx_2 \cdots \int_{-\infty}^{\infty} dx_N \exp\left(-\sum_{i=1}^N \lambda_i x_i^2\right) &= \int_{-\infty}^{\infty} dx_1 \int_{-\infty}^{\infty} dx_2 \cdots \int_{-\infty}^{\infty} dx_N \prod_{i=1}^N \exp(-\lambda_i x_i^2) \\ &= \prod_{i=1}^N \int_{-\infty}^{\infty} dx_i e^{-\lambda_i x_i^2} \\ &= \prod_{i=1}^N \sqrt{\frac{\pi}{\lambda_i}} \end{aligned} \quad (20)$$

or

$$\int_{-\infty}^{\infty} dx_1 \int_{-\infty}^{\infty} dx_2 \cdots \int_{-\infty}^{\infty} dx_N \exp\left(-\sum_{i=1}^N \lambda_i x_i^2\right) = \frac{\pi^{N/2}}{\prod_{i=1}^N \lambda_i^{1/2}} \quad (21)$$

■

First and Last Name: \_\_\_\_\_

**8. Local Minimum and Exponential Functions**

Assume that the function  $e^{-g(x)}$  has a local maximum at  $x = x_1$ . What can we say about the values of the first derivative of  $g(x)$  and the second derivative of  $g(x)$  both evaluated at  $x = x_1$ ?

**Solution:** The function  $e^{-g(x)}$  is exclusively positive and decreases to zero as the argument of the exponential gets larger. Therefore, if  $e^{-g(x)}$  has a maximum, then  $g(x)$  must be at a minimum. Namely, for  $e^{-g(x)}$  having a local maximum at  $x = x_1$ , we have

$$g'(x = x_1) = 0, \quad \text{and} \quad g''(x = x_1) > 0. \quad (22)$$

■